



**BIOLOGICAL ASSESSMENT
20062 SUMMIT DRIVE BUILDING SITE,
GOREN-LIVE OAK SITE, LOT 4
THE OAKS AT TRABUCO, ORANGE COUNTY**

UTM: 11-S: 444,500 mE; 3,726,000 mN

Prepared for
PICOA, Inc.
10866 Wilshire Boulevard, 11th Floor
Los Angeles, California 90024
(310) 441-8411 – Office
(310) 475-9112 – Fax

Prepared by
Pacific Southwest Biological Services, Inc.
P.O. Box 985
National City CA 91951-0985
Telephone 619 477 5333
15 July 2021

R. Mitchel Beauchamp, M. Sc., President
Certified Wetland Delineator #1697

PSBS # W567

PSBS #W567

**BIOLOGICAL ASSESSMENT
OF THE 20062 SUMMIT DRIVE BUILDING SITE, GOREN-LIVE OAK SITE
THE OAKS AT TRABUCO**

15 July 2021

SUMMARY

A biological survey of the one-acre development area of parcel 4 at 20062 Summit Drive at The Oaks at Trabuco, Goren property off Live Oak Canyon Road revealed a diverse mix of Scrub Oak Chaparral on north-facing slopes with Sage Scrub dominated by Cismontane Nolina and Coastal Sagebrush, along with a disturbed area on south-facing slopes. Seven sensitive animals were noted on region of the site in prior site assessments: Turkey Vulture, Coopers Hawk, Red-shouldered Hawk, Red-tailed Hawk, Coastal Cactus Wren, Coastal Rufous-crowned Sparrow, and Mule Deer.

INTRODUCTION

A biological survey of the one acre building site was performed by Pacific Southwest Biological Services, Inc., (Pacific Southwest) at the request of Mr. Gilbert Leach of Goren Financial Services, Inc., Orange County, California. The purpose of the survey was to identify sensitive biological resources and constraints in the lot grading and construction of a single-family residence within the approved nine-lot subdivision.

METHODS

The botanical portion of the survey was conducted by R. Mitchel Beauchamp on 4 June 2021, from 09:30 to 10:00 with clear skies and a temperature of 74°F. The on-foot survey covered all slope aspects, and soil types of the parcel. Particular attention was given to the Sage Scrub and Chaparral. Vegetation and sensitive plant locations were delineated on a 1" = 100' topographic map.

The zoological survey was performed during the same visit to the site by Beauchamp. All habitat types present within the study area were surveyed for wildlife. The survey included a focused survey for the Coastal California Gnatcatcher (*Polioptila californica californica*). The small size of the parcel made it possible to cover the entire site during the visit. The site was also examined by a prior assessment in 1999 (PSBS 1999).

“Pishing”, a technique commonly employed to arouse the interest of passerines, was used to call out birds. Binoculars (8.5 x 44 power) were used to assist in detection and identification.

Prior biological surveys of the immediate region were examined to assess sensitive resources known from the vicinity of the site (Pacific Southwest 1989, 1990a, 1990b, 1991, 1992, 1999).


PSBS #W567

Scientific nomenclature used in this report is from the following references: vegetation, Holland (1986); flora, Hickman (1993); birds, American Ornithologists' Union (1983, 1989); reptiles and amphibians, Collins (1990); and mammals, Jameson and Peeters (1988). Wildlife habitat delineations generally follow Mayer and Laudenslayer (1988).

LOCATION


The property and development area are situated in the south half of Section 3, Range 7 West, Township 6 South of the San Bernardino Base and Meridian; USGS 7.5' Santiago Peak, California Quadrangle in southern Orange County (Figure 1). The site is accessed from the west with Live Oak Canyon Road providing the western boundary of the property. Surrounding land use is primarily rural homesites or undeveloped. The intersection of Live Oak Canyon Road and Canyon Creek Drive lies just off-site to the northwest. The building site lies at the eastern terminus of Summit Drive.

Figure 1. Regional Location of Site



PSBS #W567

Figure 2. Vicinity Location of Site




GENERAL PHYSIOGRAPHY

The property is positioned on the south side of a ridgeline at the end of Summit Drive, and consists of the lower end of a single canyon encompassed by secondary finger ridges. High elevation at the upper end of the canyon is approximately 1278 feet, with a low elevation of 1230 feet at the mouth of the canyon. The adjacent, southern parcel is developed with a single-family home, appurtenant structure landscaping and a small vineyard. No jurisdictional wetlands or permanent watercourse occur on-site. The north-facing slope is dominated by Chaparral, while the south-facing slope gives way to Sage Scrub elements.

Soils for the site are mapped as Cieneba sandy loams in the uplands with Botella clay loams along the major drainages (Wachtell 1978). The underlying geology consists of an Oligocene nonmarine formation (Rogers 1973).

PSBS #W567

Figure 3. Areal Imagery of Site's Canyon Location



BOTANICAL RESOURCES

VEGETATION

Three primary vegetation types were delineated on the site: Disturbed Habitat, Scrub Oak Chaparral, and Diegan Sage Scrub (Figure 3). The last mentioned has an unusual component of Cismontane Nolina (*Nolina cismontana*) intermixed and includes some ecotonal chaparral elements. The Disturbed Habitat occurs at the lower, western end of the parcel. In general, Scrub Oak Chaparral predominates on all north-facing slopes while Sage Scrub occurs on the drier south- and west-facing slope.

SCRUB OAK CHAPARRAL (0.65 acre)

Dense concentrations of Scrub Oak (*Quercus berberidifolia*) are clustered on north-facing slopes where water resources tend to be greater through reduced insolation and evaporation. Interspersed with these shrubby oaks is Mountain-Mahogany (*Cercocarpus betuloides*), easily

PSBS #W567

identified during summer by its "feathered achenes," Holly-leaf Redberry (*Rhamnus ilicifolia*), Monkey (*Diplacus aurantiacus*) in the more open locales, Toyon (*Heteromeles arbutifolia*), and Chamise (*Adenostoma fasciculatum*), which occurs in larger numbers near the ridges. Erosion at the upper elevations facilitates Sage Scrub as pioneering elements over Chaparral; at numerous locales an ecotonal habitat prevails.

In many areas the Chaparral has reached a "climax stage," or senescence, with the vegetation almost impenetrable and little evidence of prior burning. Understory in such situations is minimal, but includes occasional Indian Pink (*Silene laciniata*) with its showy red petals and Filago (*Filago californica*). Sapphire Woolly Star (*Eriastrum sapphirinum* ssp. *dasyanthum*) grows in a few barren openings. Quality of the Chaparral on-site is of a standard typical of similar habitat throughout the Santa Ana foothills.

DIEGAN COASTAL SAGE SCRUB (0.44 acre)

The constituents of the Sage Scrub are primarily Coastal Sagebrush (*Artemisia californica*), Mohave Yucca (*Hesperoyucca whipplei*), and as an unusual regional element, Cismontane Nolina. This Nolina occurs in some of the highest concentrations known for this species in the Live Oak Road area. Elsewhere in southern California it is usually quite uncommon.

Coast Prickly Pear (*Opuntia occidentalis*) occurs in sporadic patches with Coast Honeysuckle (*Lonicera subspicata*) scrambling over low-growing shrubs in some areas and Bindweed (*Calystegia macrostegia* ssp. *arida*) mimicking this growth pattern over low herbaceous shrubs and annuals. Among the latter were found Stinging Lupine (*Lupinus hirsutissimus*), Cobweb Thistle (*Cirsium occidentale*), and Parry Phacelia (*Phacelia parryi*); all are beginning to decompose following an early spring bloom.

Bulbous perennials such as Golden Stars (*Bloomeria crocea*) with its bright umbel of yellow flowers, Santa Ana Mountain Mariposa Lily (*Calochortus weedii* var. *intermedius*) with its striking purplish/white petals, and Soap Plant (*Chlorogalum pomeridianum*) are uncommon on-site and typically bloom later than the annual elements.

Subshrubs in the understory include the pungent Odora (*Porophyllum gracile*), a diminutive form of Coast Goldenbush (*Isocoma menziesii* ssp. *vernonioides*) with a dense growth of branches and leaves, as well as Peony (*Paeonia californica*).

An interesting floristic addition is the presence of Turkish Rugging (*Chorizanthe staticoides* ssp. *staticoides*) with its clustered involucre and bright pink flowers. This entity is a smaller form in its floristic parts than subspecies *chrysacantha*, listed as sensitive for coastal Orange County.


Typically, the Sage Scrub on-site is of relatively high habitat quality, and is more diverse than seen to the southwest in areas such as Chiquita Canyon.

PSBS #W567

DISTURBED (0.02 acre)

A very limited area of disturbance occurs about the entrance to the canyon building site. This area is largely covered by a wood chip mulch for erosion control purposes and is primarily bereft of native vegetation.

Figure 4. Vegetation and Nolina Locations about Project Site



Flora

One hundred six species of plants were noted on the entire property in 1999, of which twenty-four are non-native, invasive elements. The current listing, with less site habitat diversity, is indicated in **Bold** font on the original listing as Appendix 1. As mentioned, the Chaparral is typical of the region, and the Sage Scrub is considered of high quality. The density of the Cismontane Nolina is a significant factor in this estimation of quality of the Sage Scrub. This species has a high density in the open space areas of the development.

PSBS #W567

Initially, floras of the region listed this species as *Nolina parryi*, still an appropriate name for plants growing in the desert foothills of Southern California. Plants in the Peninsular Ranges now described as a separate species of a more limited range and are best referred to as *Nolina cismontana*. Approximately 7 flowering specimens of the *Nolina* occur in Sage Scrub habitat of the area to be developed.

ZOOLOGICAL RESOURCES

GENERAL WILDLIFE HABITAT

Two wildlife habitats are found on this building site in the Goren property: Scrub Oak Chaparral and Diegan Sage Scrub.

SCRUB OAK CHAPARRAL

Mixed Chaparral covers a majority of the south slope of the site and contains a strong Scrub Oak and Toyon component. Canopy coverage is very dense and of moderate height, with access generally limited to the dirt roads. Rock outcrops are not present; consequently, open pockets within the vegetation are limited to dirt roads and areas of steep slopes where erosion is evident. Due to the density of the foliage, birds are the more easily detected vertebrates, with characteristic species being the Western Scrub Jay (*Aphelocoma coerulescens*), Bewick's Wren (*Thryomanes bewickii*), California Thrasher (*Toxostoma redivivum*), Wrentit (*Chamaea fasciata*), and Spotted Towhee (*Pipilo maculatus*).

DIEGAN COASTAL SAGE SCRUB

This vegetation is composed primarily of shrubs such as Coastal Sagebrush, Flat-top Buckwheat (*Eriogonum fasciculatum*), and Cismontane *Nolina* mixed with occasional Chaparral elements and annual grasses. These areas display a lower profile canopy and often a more open ground coverage. In general, such habitat is utilized by species also found in Chaparral vegetation. The greater amount of sunlight penetration enhances such areas for basking by reptiles. Two noted regional sensitive bird species are sometimes specifically associated with scrub locales as opposed to Chaparral. These are the Coastal California Gnatcatcher (*Polioptila californica californica*) and Cactus Wren (*Campylorhynchus brunneicapillus*).

Most of the on-site scrub lacks features characteristic of Gnatcatcher habitat (lack of extensive Coastal Sagebrush stands, dominance of *Nolina*, etc.), and the site's elevation generally exceeds that consistent with Gnatcatcher occurrence in Orange County (Atwood 1990). Vegetation marginally suited for the Gnatcatcher is found along a south-facing slope where a small stand of Coastal Sagebrush occurs. Detracting from its suitability is that the density of shrubs at this locale perhaps exceeds that preferred by the species, and the general density of surrounding Chaparral vegetation does not enhance the possibility of Gnatcatcher utilization.

PSBS #W567

Scattered stands of Prickly Pear Cactus are moderately suited to the presence of the Cactus Wren, and this species has been identified on adjacent parcels to the north (Pacific Southwest 1990b, 1991). On-site habitat is of comparable quality of that to the north where this species has been recently detected, and in fact the Cactus Wren was observed in each of the two major canyons on the site during the survey.

AMPHIBIANS

No amphibians were observed. Species which may occur on-site include the Pacific Chorus Frog (*Pseudacris regilla*), which may inhabit either the woodlands or dense areas of Chaparral, as well as other habitats. Both the Black-bellied Slender Salamander (*Batrachoseps nigriventris*) and Garden Slender Salamander (*Batrachoseps major*) are known from the region and could occur in any of the on-site habitats. Also, the Ensatina (*Ensatina eschscholtzi*) is frequently found living within the stick nests of the Dusky-footed Woodrat (*Neotoma fuscipes*), which were noted beneath large Chaparral shrubbery.

REPTILES

The Western Fence Lizard (*Sceloporus occidentalis*) was the only reptile observed during the survey (Appendix 2); however, the Side-blotched Lizard (*Uta stansburiana*) was noted in adjacent habitat (Pacific Southwest 1990b, 1991). Other relatively common lizards of the region which may occur on-site are the Western Skink (*Eumeces skiltonianus*) and Southern Alligator Lizard (*Elgaria multicarinata*). The San Diego Horned Lizard (*Phrynosoma coronatum blainvillei*) may also reside on-site but was not observed despite suitable season and weather conditions and the presence of its principal prey item, the Harvester Ant.

During the present site assessment, the Western Whiptail (*Cnemidophorus tigris*), was observed in the sandy canyon bottom.

The San Diego Horned Lizard was not noted during any surveys of the site or of adjacent, similar quality habitats.

Several regionally common snake species may inhabit the site or occasionally move through the property. These include the Gopher Snake (*Pituophis catenifer*), Common Kingsnake (*Lampropeltis getulus*), California Striped Racer (*Masticophis lateralis*), Coachwhip (*Masticophis flagellum*), and Southern Pacific Rattlesnake (*Crotalus viridis helleri*).

BIRDS

Thirty-six species of birds were observed on the property in 1999, with eleven additional species having been observed on-site in the immediate vicinity. Several additional species are expected to occur as winter visitors to the site. More common among these are the Hermit Thrush (*Catharus guttatus*), Ruby-crowned Kinglet (*Regulus calendula*), and Dark-eyed Junco (*Junco hyemalis*). Others, such as the Western Kingbird (*Tyrannus verticalis*), may occur as summer breeding species. Those avian species observed in the June 2021 assessment are indicated in Appendix 2.

PSBS #W567

Typical scrub/chaparral species which were detected were the California Quail (*Callipepla californica*), Anna's Hummingbird (*Calypte anna*), Western Scrub-Jay, Bushtit (*Psaltriparus minimus*), Bewick's Wren, Wrentit, California Thrasher, Phainopepla (*Phainopepla nitens*), California Towhee (*Pipilo crissalis*), Spotted Towhee, and House Finch (*Carpodacus mexicanus*). Noted on nearby parcels and expected on-site are the Greater Roadrunner (*Geococcyx californianus*), Costa's Hummingbird (*Calypte costae*), and Northern Mockingbird (*Mimus polyglottos*). The Fox Sparrow (*Passerella iliaca*) is a possible winter visitor to this habitat.

The Red-tailed Hawk (*Buteo jamaicensis*), Red-shouldered Hawk (*Buteo lineatus*), and Cooper's Hawk were the raptors observed on the site. The American Kestrel (*Falco sparverius*) has been noted in the immediate vicinity (Pacific Southwest 1990b) and is expected to occasionally forage on the subject property. The Golden Eagle (*Aquila chrysaetos*) is known from the southern Santa Ana Mountains and could forage over the site; however, the dense Chaparral and oaks offer poor foraging potential for this large bird. Also noted flying over the site was the Turkey Vulture (*Cathartes aura*), which may nest in rugged mountainous areas of rock outcrop to the north.

MAMMALS

Six species of mammals were detected on the site and three additional species were noted on nearby property (Appendix 2). Droppings of the Brush Rabbit (*Sylvilagus bachmani*) were common throughout the site and burrows of Botta's Pocket Gopher (*Thomomys bottae*) were most common in softer soils of scrub or woodlands. California Ground Squirrels (*Spermophilus beecheyi*) were observed around rural residences and pastures nearby, but were not detected on-site. Conspicuous stick nests of the Dusky-footed Woodrat were noted in dense Chaparral. Several additional small rodent species are expected to be present on-site; however, their detection generally requires a nocturnal trapping program. As none of these expected species are considered sensitive, a trapping regime was not conducted.

Larger mammals detected included two predators, the Coyote (*Canis latrans*) and the Gray Fox (*Urocyon cinereoargenteus*). Also expected in the area are the Striped Skunk (*Mephitis mephitis*), Virginia Opossum (*Didelphis virginiana*), and possibly Raccoon (*Procyon lotor*). Mule Deer (*Odocoileus hemionus*) tracks were only infrequently observed but were more plentiful on off-site trails to the north and are regularly viewed by area residents (Pacific Southwest 1991). The Bobcat (*Lynx rufus*) was noted on the properties to the north (Pacific Southwest 1990b, 1991) but no sign of its presence was observed on the subject parcel. The species is nonetheless likely to occasionally move through the property. The most formidable predator of the region is the Mountain Lion (*Felis concolor*), which has been observed by local residents (Pacific Southwest 1991). The subject property is unlikely to serve as a major use area due to the relatively low use of the site by Mule Deer, or as an important movement corridor due to the lack of a defined trail or riparian corridor connecting through to other major blocks of wild lands.

PSBS #W567

SENSITIVE PLANTS

The only sensitive plant identified on the building site was the Cismontane Nolina (*Nolina cismontana*). Several additional species are known from the region; however, the site lacks features consistent with their presence. Vernal pool species, such as Thread-leaved Brodiaea (*Brodiaea filifolia*) lack appropriate habitat on the Goren site. Sticky Dudleya (*Dudleya viscida*) occurs on metavolcanic rock in steep canyon locales such as at San Juan Hot Springs, but it lacks appropriate substrate on-site and has never been recorded this far north. Laguna Beach Dudleya (*Dudleya stolonifera*), seen in the Laguna Hills, is only known from a localized proximity to the coast and has not been collected this far inland. Many-stemmed Dudleya (*Dudleya multicaulis*) has been collected on the Starr Ranch and has the potential for occurring on-site; it grows from a small corm and is usually only detectable during late spring when in flower.

A number of sensitive species occur at higher elevations on Santiago Peak in the Santa Ana Mountains and are not expected to occur in the foothills. These include Braunton's Milkvetch (*Astragalus brauntonii*), and Santiago Peak Phacelia (*Phacelia suaveolens* ssp. *keckii*). Catalina Mariposa Lily (*Calochortus catalinae*) prefers grassland locales not well developed on the property. Prostrate Spineflower (*Chorizanthe procumbens* var. *albiflora*) has been recently collected in the region, but was not noted at the Goren site. Recently, a new taxonomic treatment of the genus has merged this entity into synonymy with other varieties, lowering its presumed sensitivity from a botanical standpoint.

SENSITIVE VERTEBRATES

Seven sensitive vertebrate species were observed on or flying over the property in 1999 but were not observed on the site during the 2021 assessment. These are Turkey Vulture (*Cathartes aura*), Golden Eagle (*Aquila chrysaetos*), Cooper's Hawk (*Accipiter cooperi*), Red-shouldered Hawk (*Buteo lineatus*), Sharp-shinned Hawk (*Accipiter striatus*), Red-tailed Hawk (*Buteo jamaicensis*), Cactus Wren (*Campylorhynchus brunneicapillus couesi*) - coastal population, Coastal Rufous-crowned Sparrow (*Aimophila ruficeps canescens*), San Diego Horned Lizard (*Phrynosoma coronatum blainvillei*), and Mountain Lion (*Felis concolor*).

EXPECTED BIOLOGICAL IMPACTS TABLE 1.SUMMARY OF IMPACTS

VEGETATION TYPE	PROPOSED DEVELOPMENT AREA ACREAGE	EXPECTED LOSS (ACRES)	REMAINING IN RESOURCE PRESERVATION AREA
Chaparral	0.65	0.65	0.7
Diegan Coastal Sage Scrub	0.44	0.44	3
Disturbed Area	0.02	0.02	0.2
TOTAL	1.11	1.11	0

Figure 5. Proposed Grading for Site Development

PSBS #W567




Figure 6. Design of proposed Single-Family Residence



Significant Impacts

The loss of 0.44 acre of Diegan Sage Scrub is considered a significant impact due to the regionally declining nature of the vegetation, and its potential to support such species as the Coastal California Gnatcatcher and Coastal Cactus Wren. Habitat of marginal or better suitability is present. Since the 1991 survey, the Coastal California Gnatcatcher has become listed as a federally Threatened Species, and impacts to Diegan Sage Scrub are generally viewed

more harshly than before as this habitat becomes a focus of a requisite Habitat Conservation Plan for the species.

Approximately 10% of the sensitive *Nolina cismontana* population will be directly impacted by grading and fill for proposed pad.

Non-Significant Impacts

The loss of 0.44 acre of good quality Scrub Oak Chaparral vegetation is not considered significant due to the absence of sensitive species associated with this habitat. Although this loss is considered an incremental impact at this time, it is noteworthy in light of the regional developments underway and proposed in the Trabuco Hills area. It should also be noted that such incremental losses lead to further habitat fragmentation and decreased value to wildlife both on-site and in surrounding areas.

The site does not contain wetland vegetation and lacks bed and bank characteristics. Due to this lack of jurisdiction, no U.S. Army Corps of Engineers 404 permit, California Department of Fish and Wildlife Section 1603 Streambed Alteration Permit, nor Regional Water Quality Control Board 401 certification are required.

RECOMMENDATIONS TO REDUCE BIOLOGICAL IMPACTS

The recommendations concerning the mitigation of impacts on the entire, multi-lot project in the preceding report have been largely implemented in the present design of this parcel and the open space allocation.

1. The vast majority of lands located outside of pad sites and fill slopes have been placed into Open Space Scenic Preservation Easements to ensure protection of these areas from clearing of native vegetation and future development. This incorporates concentrations of Diegan Coastal Sage Scrub and the heavily eroded eastern ridgeline. A 50-foot buffer of vegetation clearing to reduce fire-fuel loading would be allowed from home sites and associated structures. Selected removal is to be done near oak trees and *Nolina* colonies or individual plants.
2. Although preservation easements may include uses such as hiking and horseback riding trails by local residents, domestic pets should be maintained on leashes or restricted altogether from these areas. Hunting should be prohibited in preservation easements. Similarly, no grazing of horses or livestock should be allowed in these areas.
3. While direct impacts to *Nolina cismontana* are estimated at no greater than 15-20% of the population in this canyon, additional precautions could preclude secondary or future impacts. A fuel modification plan within the undeveloped native vegetation should have a provision that the easily-identifiable Nolinias not be grubbed or removed (they lack spine-tipped leaves). The Open Space Scenic Preservation Easement is assumed to include all native

vegetation remaining on the site. Nolinias in the area to be graded are to be salvaged for replanting on-site or offered for horticultural use.

References

- American Ornithologists' Union. 2008. Forty-ninth Supplement to the American Ornithologists' Union *Check-list of North American Birds*. *Auk* 125(3): 758-768.
- American Ornithologists' Union. 1998. Checklist of North American Birds, 7th Edition. American Ornithologists' Union. 829 pp.
- American Ornithologists' Union. 1983. Checklist of North American Birds, 6th Edition. American Ornithologists' Union.
- American Ornithologists' Union. 1989. Thirty-seventh Supplement to the American Ornithologists' Union Checklist of North American Birds. *Auk* 106: 532-538.
- Ashton, R. E., Jr. 1976. Endangered and Threatened Amphibians and Reptiles in the United States. Soc. for the Study of Amphibians and Reptiles, Herpetology Circular No. 5.
- Atwood, J. L. 1990. Status Review of the California Gnatcatcher (*Poliophtila californica*). Unpublished Technical Report, Manomet Bird Observatory, Manomet, Massachusetts. 79 pp.
- Bury, B. 1971. Status Report on California's Threatened Amphibians and Reptiles. California Department of Fish and Game, Inland Fisheries Administrative Report No. 72-2.
- California Department of Fish and Game. 1990. Special Animals. April 1990.
- California Department of Fish and Game. 1991. State and Federal Endangered and Threatened Animals of California. Revised April 1991.
- . California Department of Fish and Game. 2006. California Natural Diversity Data Base. State and Federally Listed Endangered and Threatened Animals of California. August 2006.
- Collins, Joseph T. 1990. Standard Common and Current Scientific Names for North American Amphibians and Reptiles (3rd ed.). The Society for the Study of Amphibians and Reptiles. Herpetological Circular No. 19
- Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES). 1976.
- Environmental and Energy Services Company (ERCE). 1989. Phase I Report Amber Ridge California Gnatcatcher Study. Prepared for Weingarten, Siegel, Fletcher Group, Inc. 15 November 1989

- Hamilton, R. A. and D. R. Willick. 1996. The birds of Orange County, California: Status and distribution. Sea & Sage Press, Sea & Sage Audubon Society; 1 edition. 150 pp.
- Hickman, J. P. A. 1993. Jepson Manual of the Higher Plants of California. University of California Press, Berkeley. 1400pp.
- Holland, Robert F. 1986. Preliminary Descriptions of the Terrestrial Natural Communities of California. California Department of Fish and Game.
- International Union for the Conservation of Nature and Natural Resources. 1979. Red Data Book, Vol. 3: Amphibia and Reptilia.
- Jameson, E.W., Jr. and H. J. Peeters. 2004. California Mammals. University of California Press, Berkeley. 440 pp.
- Mayer, Kenneth E. and William F. Laudenslayer, Jr., editors. 1988. A Guide to Wildlife Habitats of California. California Department of Forestry and Fire Protection.
- Oberbauer, T. 1996. Terrestrial Vegetation Communities in San Diego County based on Holland's Descriptions. San Diego Association of Governments, San Diego, California. 6 pp.
- Oberbauer, T. 2005. Terrestrial Vegetation Communities in San Diego County based on Holland's Descriptions. San Diego Association of Governments, San Diego, California. Revised March 2005. 5 pp.
- Pacific Southwest Biological Services, Inc. 1989. Wildlife Dispersal Study, Hunt Lodge Relocation Study Area, Coto De Caza, Orange County. Prepared for Coto De Caza Development Corporation. 12 December 1989.
- Pacific Southwest Biological Services, Inc. 1990a. Report of a Biological Survey and Assessment of the Proposed Bridlewood Development Site, Trabuco Area, Orange County, California. Prepared for R.B. McComic Inc., Laguna Niguel, California. 2 October 1990.
- Pacific Southwest Biological Services, Inc. 1990b. Report of a Biological Assessment of Live Oak Estates, Orange County, California. Prepared for Aston Group of Santa Ana, Santa Ana, California. 5 October 1990.
- Pacific Southwest Biological Services, Inc. 1991. Report of a Biological Assessment of the 20 Acre Gooding/La Prairie Site. Prepared for Land Planning/Governmental Relations, Irvine, California. 18 June 1991.
- Pacific Southwest Biological Services. 1999. Biological Assessment of the 34.4 acre Goren-Live Oak Site, The Oaks at Trabuco. 30 Sept. 1999.

PSBS #W567

- Rea, Amadeo M., and Kenneth L. Weaver. 1990. The Taxonomy, Distribution, and Status of Coastal California Cactus Wrens. **Western Birds** 21: 80-126.
- Remsen, J. V., II. 1980. Bird Species of Special Concern in California. California Department of Fish and Game, Sacramento, California. 54pp.
- Rogers, Thomas H. 1973. Geologic Map of California, Santa Ana Sheet. California Division of Mines and Geology, Sacramento, California.
- Stewart, G. R. 1971. Rare, Endangered and Depleted Amphibians and Reptiles in California. **Herpetology** 5: 29-35.
- Tate, James, Jr. 1986. The Blue List for 1986. **American Birds** 40(2):227-236.
- U.S. Fish and Wildlife Service. 1986. Endangered and Threatened Wildlife and Plants. Code of Fed. Regul. Title 50, Part 17.11 and 17.12 (revised January 1, 1986).
- U.S. Fish and Wildlife Service. 1989. Endangered and Threatened Wildlife and Plants. **Federal Register** Part IV, January 6, 1989.
- U. S. Fish and Wildlife Service. 1997. Coastal California Gnatcatcher (*Poliophtila californica californica*) Presence/Absence Survey Guidelines, February 28, 1997. 4 pp.
- U.S. Fish and Wildlife Service. 1999. Bradley, J. As quoted in San Diego *Union Tribune*. March 11, 1999.
- Wachtell, John K. 1978. Soil Survey of Orange County and Western Part of Riverside County, California. United States Department of Agriculture and Soil Conservation Service and Forest Service, in Cooperation with University of California.

Appendix 1. Floral Checklist of the Goren-Trabuco Oaks Site.

Bold entries indicate referenced building site

HABITAT C = Scrub Oak Chaparral W = Coast Live Oak Riparian Woodland

D = Diegan Sage Scrub

CRYPTOGAMS Habitats

SPIKE-MOSSES

Selaginellaceae-Spike-Moss Family

Selaginella bigelovii Underw. Common Spike-Moss C

GYMNOSPERMS

Pinaceae

**Pinus halepensis* Aleppo Pine D

DICOTYLEDONS

Aizoaceae - Carpet-weed Family

**Aptenia cordifolia* (L.) Schwant W

Anacardiaceae - Sumac Family

Malosma laurina (Nutt.) Nutt. ex Abrams Laurel-Leaf Sumac D

***Rhus integrifolia* (Nutt.) Benth. & Hook. Lemonade Berry D**

***Toxicodendron radicans* ssp. *diversilobum* (T. & G.) Thorne Poison-Oak W**

Apiaceae - Carrot Family

Apiastrum angustifolium Nutt. Mock Parsley W

Daucus pusillus Michx. Rattlesnake Weed D

**Foeniculum vulgare* Mill. Fennel W

***Yabea macrocarpa* (Hook. & Arn.) Koso-Polj. C**

Asclepiadaceae - Milkweed Family

Asclepias eriocarpa Benth. Indian Milkweed W

Asteraceae - Sunflower Family

**Lactuca serriola* L. Wild Lettuce C

Acourtia microcephala DC. Sacapellote, Purpleheads C

Ambrosia psilostachya DC. var. *californica* (Rydb.) Blake Western Ragweed W

***Artemisia californica* Less. California Sagebrush D**

Artemisia douglasiana Bess. in Hook. Mugwort W

Artemisia dracuncululus L. Dragon Sagewort W

Baccharis pilularis ssp. *consanguinea* (DC.) C.B. Wolf Coyote Brush W

***Baccharis salicifolia* (R. & P.) Pers. Mule-fat W**

* *Carduus pycnocephalus* L. Italian Thistle C, W

* *Centaurea melitensis* L. Tocalote D

Chaenactis artemisiaefolia (Harv. & Gray) Gray Pincushion D

Cirsium occidentale (Nutt.) Jeps. Cobweb Thistle D

Encelia californica Nutt. California Encelia D

Erigeron foliosus var. *foliosus* Leaf Daisy D

***Eriophyllum confertiflorum* (DC.) Gray var. *confertiflorum* Golden-Yarrow C,D**

Filago californica Nutt. California Filago C,D

Appendix 1. Floral Checklist of the Goren-Trabuco Oaks Site (continued)

Asteraceae- Daisy Family (continued)

* <i>Filago gallica</i> L. Narrow-leaf Filago	D	
<i>Gnaphalium beneolens</i> Davids. Fragrant Everlasting	D	
<i>Gnaphalium (Pseudognaphalium) californicum</i> DC. California Everlasting	C	
<i>Gnaphalium microcephalum</i> Nutt. White Everlasting	D	
<i>Hazardia squarrosa</i> ssp. <i>grindelioides</i> (DC.)Clarke Sawtooth Goldenbush	D,S	
* <i>Hedypnois cretica</i> (L.) Willd. Crete Hedypnois	D	
<i>Hemizonia fasciculata</i> (DC.) Torrey & Gray Fascicled Tarplant	D	
<i>Heterotheca grandiflora</i> Nutt. Telegraph Weed	D	
* <i>Hypochoeris glabra</i> L. Smooth Cat's-ear	D	
* <i>Lactuca serriola</i> L. Wild Lettuce	D	
<i>Isocoma menziesii</i> var. <i>vernonioides</i> (Nutt.)Nesom Coastal Goldenbush	D	
<i>Lessingia filaginifolia</i> (Hook. & Arn.) M.A. Lane var. <i>filaginifolia</i> Cudweed Aster	C	
<i>Porophyllum gracile</i> Benth. Odora	D	
<i>Solidago californica</i> Nutt. California Goldenrod	W	
*<i>Sonchus oleraceus</i> L. Common Sow-Thistle	D	
<i>Stebbinoseris heterocarpa</i> (Nutt.) Chambers	C	
<i>Stephanomeria diegensis</i> Gottlieb San Diego Wreath-Plant	D	
<i>Stephanomeria virgata</i> Benth. ssp. <i>virgata</i> Virgate Wreath-plant	D	
<i>Stylocline gnaphalioides</i> Nutt. Everlasting Nest Straw	D	
Boraginaceae - Borage Family		
<i>Cryptantha intermedia</i> (Gray)Greene Nievitas	D	
Brassicaceae - Mustard Family		
*<i>Hirschfeldia incana</i> (L.) Lagr.-Fossat Short-pod Mustard	D	
* <i>Sisymbrium irio</i> L. London Rocket	D	
Cactaceae - Cactus Family		
<i>Opuntia littoralis</i> (Engelm.)Ckll. var. <i>littoralis</i> Coast Prickly-Pear	D	
<i>Opuntia occidentalis</i> Engelm. & Bigel. Western Prickly-Pear, Thicket Tuna	D	
<i>Opuntia vaseyi</i> (J. Coulter) Britt & Rose	D	
Caprifoliaceae - Honeysuckle Family		
<i>Lonicera subspicata</i> var. <i>denudata</i> Rehd. San Diego Honeysuckle	D	
Caryophyllaceae - Pink Family		
<i>Silene laciniata</i> ssp. <i>major</i> Hitchc. & Maguire Southern Pink	D	
Chenopodiaceae - Goosefoot Family		
* <i>Chenopodium ambrosioides</i> L. Mexican-Tea	W	
Convolvulaceae - Morning-Glory Family		
<i>Calystegia macrostegia</i> ssp. <i>arida</i> (Greene)Brumm. Finger-leaf Morning-Glory	D	
<i>Cuscuta californica</i> H. & A. Witch's Hair	D	
Crassulaceae - Stonecrop Family		
<i>Dudleya pulverulenta</i> (Nutt.)Britt. & Rose Chalk-lettuce	D	
Cucurbitaceae - Gourd Family		
<i>Marah macrocarpus</i> (Greene)Greene Manroot, Wild-Cucumber	C	

Appendix 1. Floral Checklist of the Goren-Trabuco Oaks Site (continued)

Euphorbiaceae - Spurge Family	
<i>Eremocarpus setigerus</i> (Hook.)Benth. Doveweed	D
Fabaceae - Pea Family	
<i>Lathyrus laetiflorus</i> ssp. <i>alefeldii</i> (White)Brads. Sweetpea	D
<i>Lotus purshianus</i> (Benth.)Clem. & Clem. Spanish-Clover	D
<i>Lotus scoparius</i> (Nutt. in T. & G.)Ottley ssp. <i>scoparius</i> (<i>Acmispon glaber</i>) Deerweed	D
<i>Lupinus hirsutissimus</i> Benth. Stinging Lupine	D
* <i>Medicago polymorpha</i> L. California Bur-Clover	D
<i>Trifolium tridentatum</i> Lindl. var. <i>tridentatum</i> Tomcat Clover	C
Fagaceae - Oak Family	
<i>Quercus agrifolia</i> Neé var. <i>agrifolia</i> Coast Live Oak	W
<i>Quercus berberidifolia</i> Liebm. Scrub Oak	C
Geraniaceae - Geranium Family	
* <i>Erodium cicutarium</i> (L.)L'Hér. Red-stem Filaree	D
Hydrophyllaceae - Waterleaf Family	
<i>Eucrypta chrysanthemifolia</i> (Benth.) Greene var. <i>chrysanthemifolia</i>	D
<i>Phacelia parryi</i> Torr.	D
<i>Phacelia suffrutescens</i> (Parry) Const.	D
Lamiaceae - Mint Family	
<i>Salvia apiana</i> Jeps. White Sage	D
<i>Salvia mellifera</i> Greene Black Sage	C,D
Myrtaceae - Myrtle Family	
* <i>Eucalyptus</i> sp.	W
Nyctaginaceae - Four-O'Clock Family	
<i>Mirabilis californica</i> Gray Wishbone Plant	D
Oxalidaceae - Wood-Sorrel Family	
<i>Oxalis albicans</i> ssp. <i>californica</i> (Abrams)Eiten California Wood-Sorrel	D
Paeoniaceae - Peony Family	
<i>Paeonia californica</i> Nutt. in T. & G. California Peony	C
Plantaginaceae - Plantain Family	
* <i>Plantago lanceolata</i> L. English Plantain	W
Platanaceae - Sycamore Family	
<i>Platanus racemosa</i> Nutt. Western Sycamore	W
Polemoniaceae	
<i>Eriastrum sapphirinum</i> ssp. <i>dasyanthum</i> (Brand)Mason Woolly-Star	C
<i>Leptodactylon californicum</i> H. & A. Prickly Phlox	C,D
Polygonaceae - Buckwheat Family	
<i>Chorizanthe staticoides</i> Benth. ssp. <i>staticoides</i> Turkish Rugging	D
<i>Eriogonum fasciculatum</i> Benth. Flat-top Buckwheat	D
* <i>Rumex crispus</i> L. Curly Dock	W
Rhamnaceae - Buckthorn Family	
<i>Ceanothus crassifolius</i> Torr. Thick-leaf Ceanothus	C
<i>Rhamnus</i> (<i>Endotropis</i>) <i>ilicifolia</i> Kell. Hollyleaf Redberry	C

Appendix 1. Floral Checklist of the Goren-Trabuco Oaks Site (continued)

Rosaceae - Rose Family	
<i>Adenostoma fasciculatum</i> H. & A. Common Chamise	C
<i>Cercocarpus betuloides</i> Nutt. ex T. & G. Mountain-Mahogany	C
<i>Heteromeles arbutifolia</i> (Ait.) M. Roem. Hollywood, Toyon	C,D
Rubiaceae - Madder Family	
<i>Galium angustifolium</i> Nutt., T. & G. ssp. <i>angustifolium</i> Narrow-leaf BedstrawD	
Scrophulariaceae - Figwort Family	
<i>Cordylanthus rigidus</i> Nutt. ex Benth. Dark-tip Bird's-Beak	C,D
<i>Diplacus aurantiacus</i> ssp. <i>australis</i> (McMinn) R.M. Beeks ex Thorne	C
<i>Keckiella cordifolia</i> (Benth.) Straw Climbing Penstemon	C
Solanaceae - Nightshade Family	
* <i>Nicotiana glauca</i> Grah. Tree Tobacco	D
MONOCOTYLEDONS Habitat	
Agavaceae - Agave Family	
<i>Nolina cismontana</i> Dice.	D
<i>Yucca (Hesperoyucca) whipplei</i> Torr. Our Lord's Candle	C,D
Alliaceae - Onion Family	
<i>Bloomeria crocea</i> (Torr.)Cov. ssp. <i>crocea</i> Common Golden-Stars	D
<i>Dichelostemma pulchellum</i> (Salisb.) Heller Wild-Hyacinth	D
Iridaceae - Iris Family	
<i>Sisyrinchium bellum</i> Wats. Blue-eyed-Grass	D
Juncaceae - Rush Family	
<i>Juncus dubius</i> Engelm. Mariposa Rush	W
Liliaceae - Lily Family	
<i>Calochortus weedii</i> Wood var. <i>intermedius</i> Weed's Mariposa-Lily	D
<i>Chlorogalum parviflorum</i> Wats. Small-Flower Soap-Plant	D
<i>Chlorogalum pomeridianum</i> (DC.) Kunth Wavy-Leaf Soap-Plant	D
Poaceae - Grass Family	
<i>Agropyron parishii</i> var. <i>laeve</i> Scribn. & Sm. California Wheatgrass	C
* <i>Avena barbata</i> L. Slender Oat	C,D
<i>Bothriochloa barbinodis</i> (Lag.)Herter Plumed Beardgrass	D
*<i>Bromus diandrus</i> Roth. Ripgut Grass	C,D
* <i>Bromus mollis</i> L. Soft Chess	C,D
* <i>Bromus rubens</i> L. Red Brome	C,D
* <i>Cynodon dactylon</i> (L.) Pers. Bermuda Grass	D
* <i>Festuca myuros</i> L.	C,D
* <i>Hordeum murinum</i> ssp. <i>glaucum</i> (Steud.)Tzvel. Glaucous Barley	D
* <i>Lamarckia aurea</i> (L.) Moench Goldentop	D
<i>Leymus condensatus</i> (Presl) A. Love Giant Wild Rye	D,W
<i>Melica imperfecta</i> Trin. Coast Range Melic	D
C	
<i>Nassella lepida</i> (A.S. Hitchcock) Barkworth Foothill Needlegrass	D
*<i>Piptatherum miliaceum</i> (L.) Cosson. Smilo Grass	W
* <i>Polypogon monspeliensis</i> (L.) Desf. Annual Beardgrass	W
* - Denotes non-native plant taxa	

Appendix 2. Animals Observed or Detected on the Goren-Trabuco Oaks Site

Habitat: C = Coastal Scrub F = Flyover M = Mixed Chaparral O = Oak Woodland
 p = Present on adjacent property (Pacific Southwest 1990, 1991), may occur on subject property as well

Common Name	Scientific Name	Number/Means of Detection	Habitat
Reptiles			
Phrynosomatidae			
Western Fence Lizard	<i>Sceloporus occidentalis</i>		C
Side-blotched Lizard	<i>Uta stansburiana</i>		
Orange-throat Whiptail	<i>Cnemidophorus hyperythrus beldingi</i>		C
Colubridae (Colubrids)			
Gopher Snake	<i>Pituophis catenifer</i>		p
Viperidae (Vipers)			
Southern Pacific Rattlesnake	<i>Crotalus viridis helleri</i>		p
Birds			
Cathartidae (American Vultures)			
Turkey Vulture	<i>Cathartes aura</i>		F
Accipitridae (Hawks, Old World Vultures, and Harriers)			
Cooper's Hawk	<i>Accipiter cooperi</i>		M
Red-tailed Hawk	<i>Buteo jamaicensis</i>		F
Red-shouldered Hawk	<i>Buteo lineatus</i>		p
Falconidae (Caracaras and Falcons)			
American Kestrel	<i>Falco sparverius</i>		p
Phasianidae (Quails, Pheasants, and Relatives)			
California Quail	<i>Callipepla californica</i>		M
Columbidae (Pigeons and Doves)			
Mourning Dove	<i>Zenaida macroura</i>		O
Apodidae (Swifts)			
White-throated Swift	<i>Aeronautes saxatalis</i>		p
Trochilidae (Hummingbirds)			
Anna's Hummingbird	<i>Calypte anna</i>		C
Costa's Hummingbird	<i>Calypte costae</i>		p
Picidae (Woodpeckers and Wrynecks)			
Acorn Woodpecker	<i>Melanerpes formicivorus</i>		O
Nuttall's Woodpecker	<i>Picoides nuttallii</i>		O
Northern Flicker	<i>Colaptes auratus</i>		O

		Number/Means	
Appendix 2. Animals Observed or Detected on the Goren-Trabuco Oaks Site			
Common Name	Scientific Name	of Detection	Habitat
Tyrannidae (Tyrant Flycatchers)			
Pacific-slope Flycatcher	<i>Empidonax difficilis</i>		p
Ash-throated Flycatcher	<i>Myiarchus cinerascens</i>		C
Western Kingbird	<i>Tyrannus verticalis</i>		M
Hirundinidae (Swallows)			
Northern Rough-winged Swallow	<i>Stelgidopteryx serripennis</i>		F
Cliff Swallow	<i>Hirundo pyrrhonota</i>		p
Corvidae (Jays, Magpies, and Crows)			
Western Scrub Jay	<i>Aphelocoma coerulescens</i>		M,O
American Crow	<i>Corvus brachyrhynchos</i>		F
Common Raven	<i>Corvus corax</i>		F
Paridae (Titmice)			
Oak Titmouse	<i>Parus inornatus</i>		O
Aegithalidae (Bushtit)			
Bushtit	<i>Psaltriparus minimus</i>		M,O
Troglodytidae (Wrens)			
Cactus Wren	<i>Campylorhynchus brunneicapillus</i>		p
Bewick's Wren	<i>Thryomanes bewickii</i>		C,M
House Wren	<i>Troglodytes aedon</i>		O
Muscicapidae (Old World Warblers, Gnatcatchers, Kinglets, Thrushes, Bluebirds, and Wrentit)			
Wrentit	<i>Chamaea fasciata</i>		C,M,O
Mimidae (Mockingbirds and Thrashers)			
Northern Mockingbird	<i>Mimus polyglottos</i>		p
California Thrasher	<i>Toxostoma redivivum</i>		M
Ptilonotidae (Silky Flycatchers)			
Phainopepla	<i>Phainopepla nitens</i>		M
Sturnidae (Starlings)			
European Starling	<i>Sturnus vulgaris</i>		p
Vireonidae (Typical Vireos)			
Hutton's Vireo	<i>Vireo huttoni</i>		p
Emberizidae (Warblers, Sparrows, Blackbirds and Relatives)			
Coastal Rufous-crowned Sparrow	<i>Aimophila ruficeps canescens</i>		C
Song Sparrow	<i>Melospiza melodia</i>		O
White-crowned Sparrow	<i>Zonotrichia leucophrys</i>		O
Yellow-rumped Warbler	<i>Dendroica coronata</i>		O
Orange-crowned Warbler	<i>Vermivora celata</i>		O
California Towhee	<i>Pipilo crissalis</i>		C,M,O
Spotted Towhee	<i>Pipilo macalatus</i>		M,O
Fringillidae (Finches)			
House Finch	<i>Carpodacus mexicanus</i>		F,M
Lesser Goldfinch	<i>Carduelis psaltria</i>		M,O

American Goldfinch	<i>Carduelis tristis</i>	p
--------------------	--------------------------	---

Appendix 2. Animals Observed or Detected on the Goren-Trabuco Oaks Site

Common Name	Scientific Name	Number/Mean of Detection	Habitat
Mammals			
Leporidae (Rabbits and Hares)			
Brush Rabbit	<i>Sylvilagus bachmani</i>		C,M
Sciuridae (Squirrels, Chipmunks, and Marmots)			
California Ground Squirrel	<i>Spermophilus beecheyi</i>		p
Geomyidae (Pocket Gophers)			
Botta's Pocket Gopher	<i>Thomomys bottae</i>		C,O
Muridae (Rats, mice, and voles)			
Dusky-footed Woodrat	<i>Neotoma fuscipes</i>		M,O
Canidae (Foxes, Wolves, and Relatives)			
Coyote	<i>Canis latrans</i>		C,M
Gray Fox	<i>Urocyon cinereoargenteus</i>		C,M
Felidae (Cats)			
Mountain Lion	<i>Felis concolor</i>		p
Bobcat	<i>Lynx rufus</i>		p
Cervidae (Deer, Elk, and Relatives)			
Mule Deer	<i>Odocoileus hemionus</i>		M

Appendix 3. Site Photographs W567 Trabuco Oaks Lot # 7, 20062 Summit Trail



Western Disturbed area



Chaparral on S side of site



View to NE up canyon in ecotone area



Lower , W end of site



Nolinas in Sage Scrub on N side of site



View to W in Sage Scrub habitat



Male and female Nolinas on N side of site



SageScrub on N side of site



Chaparral in bottom of site



Chaparral and Sage Scrub ecotone in bottom of site



View to E of Sage Scrub and Chaparral